

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



United States
Department of
Agriculture

Agricultural
Research
Service

ARS-19

November 1984

Evaluation of New Canal Point Sugarcane Clones

1983-84 Harvest Season

SEP 24 1984

ABSTRACT

Glaz, B., J. D. Miller, P. Y. P. Tai, J. L. Dean, M. S. Kang, and O. Sosa, Jr. 1984. Evaluation of New Canal Point Sugarcane Clones: 1983-84 Harvest Season. U.S. Department of Agriculture, Agricultural Research Service, ARS-19, 24 p.

Thirty-two new Canal Point (CP) clones of sugarcane (11 in plant cane, 10 in first-ratoon cane, and 11 in second-ratoon cane) were grown in replicated experiments and harvested at 9 locations representing 5 soil series (Terra Ceia muck, Pahokee muck, Lauderdale muck, Torry muck, and Pompano fine sand). The cane and sugar yields of these clones, interspecific hybrids of Saccharum spp., were compared with those of CP 70-1133 in plant-cane and first-ratoon experiments and CP 63-588 in second-ratoon experiments. Each clone was also rated for its reaction to sugarcane smut, Ustilago scitaminea, in separate inoculation tests and by natural infection and for its reaction to sugarcane rust, Puccinia melanocephala, by natural infection. There were no clones superior to CP 70-1133 in the plant-cane experiments. In the first-ratoon experiments, CP 78-1247 was the only clone with a significantly higher yield, in metric tons of sugar per hectare (TS/H), than CP 70-1133. Other promising clones from the first-ratoon experiments were CP 78-1140, CP 78-1156, CP 78-1263, CP 78-1610, CP 78-1628, and CP 78-2114, all of which were not significantly different from CP 70-1133 in yield of TS/H. In the second-ratoon experiments, CP 77-1414 was the only clone with adequate disease resistance for commercial production which had a significantly higher yield of TS/H than CP 63-588. CP 77-1776, a clone which had high yields of sugar (in kilograms) per metric ton of cane (KS/T) in the two previous harvest seasons, was once again significantly higher than CP 63-588 in this characteristic, both at early and regular harvests. However, CP 77-1776 was lower than CP 63-588 in yield estimates of TS/H from early samples and at harvest time.

Keywords: Florida, Lauderdale muck, Pahokee muck, Pompano fine sand, Puccinia melanocephala, Saccharum spp., sugarcane cultivars, sugarcane rust, sugarcane smut, sugarcane varieties, sugarcane yields, sugar yields, Terra Ceia muck, Torry muck, Ustilago scitaminea.

CONTENTS

| | |
|------------------------------|---|
| Introduction | 1 |
| Test procedures | 2 |
| Results and discussion | 3 |
| Plant cane | 3 |
| First-ratoon cane | 3 |
| Second-ratoon cane | 4 |
| Summary | 5 |
| References | 6 |

TABLES

| | |
|---|----|
| 1. Parentage, variety correction factors, and ratings for smut and rust susceptibility of CP 63-588, CP 70-1133, and 32 new sugarcane clones | 7 |
| 2. Yields of cane, in metric tons per hectare, from plant cane on Lauderdale, Pahokee and Torry muck and on Pompano fine sand | 8 |
| 3. Indicated yields of 96° sugar, in kilograms per metric ton of cane, from preharvest samples of plant cane on Lauderdale, Pahokee, and Torry muck and on Pompano fine sand | 9 |
| 4. Indicated yields of 96° sugar, in metric tons per hectare, from preharvest samples of plant cane on Lauderdale, Pahokee, and Torry muck and on Pompano fine sand | 10 |
| 5. Indicated yields of 96° sugar, in kilograms per metric ton of cane, from plant cane on Lauderdale, Pahokee, and Torry muck and on Pompano fine sand | 11 |
| 6. Indicated yields of 96° sugar, in metric tons per hectare, from plant cane on Lauderdale, Pahokee, and Torry muck and on Pompano fine sand | 12 |
| 7. Yields of cane, in metric tons per hectare, from first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand | 13 |
| 8. Indicated yields of 96° sugar, in kilograms per metric ton of cane, from preharvest samples of first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand | 14 |
| 9. Indicated yields of 96° sugar, in metric tons per hectare, from preharvest samples of first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand | 15 |
| 10. Indicated yields of 96° sugar, in kilograms per metric ton of cane, from first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand | 16 |

| | | |
|-----|--|----|
| 11. | Indicated yields of 96° sugar, in metric tons per hectare, from first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand ... | 17 |
| 12. | Yields of cane, in metric tons per hectare, from second-ratoon cane on Lauderdale, Terra Ceia, and Torry muck and on Pompano fine sand | 18 |
| 13. | Indicated yields of 96° sugar, in kilograms per metric ton of cane, from preharvest samples of second-ratoon cane on Lauderdale, Terra Ceia, and Torry muck and on Pompano fine sand | 19 |
| 14. | Indicated yields of 96° sugar, in metric tons per hectare, from preharvest samples of second-ratoon cane on Lauderdale, Terra Ceia, and Torry muck and on Pompano fine sand | 20 |
| 15. | Indicated yields of 96° sugar, in kilograms per metric ton of cane, from second-ratoon cane on Lauderdale, Terra Ceia, and Torry muck and on Pompano fine sand ... | 21 |
| 16. | Indicated yields of 96° sugar, in metric tons per hectare, from second-ratoon cane on Lauderdale, Terra Ceia, and Torry muck and on Pompano fine sand | 22 |

ACKNOWLEDGMENTS

The authors are grateful to the Florida Sugar Cane League, Inc., for making available John Adams to assist in planting, sampling, harvesting, and analyzing, and Howard Weir to assist in harvesting the field experiments described in this report; and Betty Murphy to type the manuscript.

This publication is available from the U.S. Sugarcane Field Station, Star Route, Box 8, Canal Point, FL 33438, while supply lasts.

Copies may also be purchased from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.

EVALUATION OF NEW CANAL POINT
SUGARCANE CLONES: 1983-84
Harvest Season

By Glaz, B., J. D. Miller, P. Y. P. Tai,
J. L. Dean, M. S. Kang, and O. Sosa, Jr. ¹

INTRODUCTION

Clonal selection at precommercial stages is one of the major components involved in the successful production of sugarcane, interspecific hybrids of Saccharum spp. Although production of sugar per unit area is a very important characteristic, it is not the only yield factor upon which sugarcane is evaluated. In addition, analyses are made on the quantity of cane needed to produce a particular sugar yield and on the relative millability of the cane. The time of year and the amount of time during the year that a clone yields its highest amount of sugar per unit area can be very important, since sugarcane harvest seasons are usually long. Sugarcane is also characteristically grown as plant and ratoon crops; thus, clones are evaluated accordingly.

Clones that have the desired agronomic characteristics must also be productive in the presence of major disease, insect, and weed pests. Determination of pest resistance and tolerance can be complicated processes that take several years to complete. The selection team must be careful not to discard clones that could be grown commercially because of sufficient resistance or tolerance to pests, but it must also be careful to discard clones that are too susceptible to pests to be grown commercially.

^{1/} Glaz is an agronomist; Miller and Tai are research geneticists; Dean is a research plant pathologist; and Sosa, research entomologist--all with the U. S. Department of Agriculture, Agricultural Research Service, U. S. Sugarcane Field Station, Star Route, Box 8, Canal Point, Florida 33438. Kang is assistant professor, Everglades Research and Education Center, Institute of Food and Agricultural Sciences, University of Florida, Belle Glade 33430.

In recent years in many sugarcane producing regions worldwide, there have been major pest attacks on widely grown commercial clones. It is now generally accepted that sugarcane growing regions need adequate reserves of clones (even though yields of these reserve clones may not be as high as those of the major commercial clones) so that major clones can be quickly replaced if they show susceptibility to a new pest or a new race of an established pest. In Florida, three important commercial clones, CP 72-1210, CL 54-378, and CP 74-2005, are threatened by what are probably at least two new races of sugarcane rust, Puccinia melanocephala H. Syd. & P. Syd.

Each year at Canal Point, approximately 100,000 seedlings are evaluated from crosses made with a diverse germplasm collection, although perhaps one with not a sufficiently diverse cytoplasmic base (Mangelsdorf 1983). The clones reported on herein are receiving their last 3 years of evaluation by us. Each clone that successfully passes the first year of this phase must undergo an additional 3 to 5 years of evaluation by Florida sugarcane industry personnel before a decision is made on whether it will become a commercial clone.

Characteristics of clones that may be valuable for breeding with other clones are identified throughout the selection process. Clones with specific characteristics are sought by sugarcane breeders worldwide. From September 1983 to May 1984, Canal Point clones were sent to Australia, Colombia, France, Iraq, Malaysia, Mauritius, Puerto Rico, and Zimbabwe. Canal Point clones also were sent to Clewiston, Gainesville, and Orlando in Florida and to Louisiana, Maryland, Ohio, and Virginia.

TEST PROCEDURES

Twenty-three replicated experiments of 32 new Canal Point (CP) clones (11 in plant cane, 10 in first-ratoon cane, and 11 in second-ratoon cane) and the reference clones (CP 63-588 and CP 70-1133) were harvested at 9 growers' farms. CP 70-1133 was the reference clone in plant-cane and first-ratoon experiments and CP 63-588 was the reference clone in second-ratoon experiments. There were plant-cane, first-ratoon, and second-ratoon experiments at each location except Okeelanta, which did not have a second-ratoon experiment; Beardsley Farm, which did not have a plant-cane experiment; and Eastgate Farms, which had only a plant-cane experiment. Ten experiments were conducted on Lauderhill muck. Of these, eight experiments were at A. Duda and Sons' Farm (east of Belle Glade), Gulf and Western Food Products Co. (Okeelanta), and A. F. Saunders, Inc. (south of Clewiston). The other two were second-ratoon experiments at New Farm, Inc. (east of Canal Point), and Wedgworth Farms, Inc. (east of Belle Glade). Four experiments were conducted on Pahokee muck, which is similar to Lauderhill and Terra Ceia muck but deeper than the former and not as deep as the latter (Snyder et al. 1978): plant-cane experiments at New Farm and Wedgworth Farms and plant-cane and first-ratoon experiments at South Florida Industries (near 20 Mile Bend in Palm Beach County). Experiments on Terra Ceia muck consisted of first-ratoon experiments at New Farm and Wedgworth Farms and a second-ratoon experiment at South Florida Industries. Torrey muck was the soil for the plant-cane experiment at Eastgate Farms (north of Belle Glade) and two ratoon experiments at Beardsley Farm (near Lake Harbor); and Pompano fine sand, for three experiments at the Lykes Brothers' Farm (near Moorehaven in Glades County).

Selection of each clone for planting in these replicated experiments was based on 5 years of data. Beginning

last year, and as reflected by results presented this year for the first-ratoon experiments, we replaced CP 63-588 with CP 70-1133 because it had become the most widely grown commercial clone in Florida (Donovan and Glaz 1983). In each of the 23 experiments, clones were planted with 2 lines of seed cane per furrow in plots of 0.0065 hectare in a randomized complete-block design with 4 replications. Each plot was 10.7 meters long and 6.1 meters wide. The distance between rows was 1.5 meters, and a distance of 1.5 meters separated the end of each plot row and the beginning of the next plot row. The margins of the experiments were buffered to reduce mechanical damage and border effects, but individual four-row plots were not buffered.

Sugarcane management practices, such as fertilizing, cultivating, pest control, burning, loading, and hauling, were the same for each experimental plot as for the commercial field in which the plot was located.

To evaluate early-season sugar production, 10 stalks per plot were randomly sampled from the unburned cane in 2 of the 4 replications at each location between October 17 and 26, 1983. These samples were milled, the crusher juice was analyzed for Brix and sucrose, and the indicated yields of sugar in kilograms per metric ton of cane were determined. To estimate the yield of sugar per hectare from these preharvest data, we assumed that the preharvest yield of cane per hectare was equal to the actual yield of cane per hectare obtained at harvest.

All experiments were harvested between October 30, 1983, and February 27, 1984. However, 19 of the 23 experiments were harvested in January and February 1984. The ranges of harvest dates were as follows: January 3 to February 27, 1984, for plant-cane; December 7, 1983 to February 24, 1984, for first-ratoon cane; and October 30, 1983, to February

RESULTS AND DISCUSSION

12, 1984, for second-ratoon cane. After each plot had been burned, all cane was cut and piled by hand and then weighed with a tractor-mounted weighing device. Fifteen stalks were randomly selected from each plot and transported to the Agricultural Research Service's laboratory at Canal Point for weighing, milling, and crusher-juice analysis.

All values for yield of sugar per metric ton of cane and for yield of sugar per hectare in this report are indicated (theoretical) yields calculated in accordance with a simplification of the Winter-Carp-Geerligs formula (Arceneaux 1935); an explanation of the formula was given previously (Rice and Hebert 1972).

Although the indicated sugar yields reported herein may not be obtained by all sugar factories, they are valid for comparing clones having different milling qualities. Varietal correction factors were used in all the theoretical sugar calculations.

Statistical analyses were done according to McIntosh (1983). F -ratios were chosen according to a mixed model, with treatments (clones) fixed and locations random. The source of variation that corresponded to the error term for the effect being tested was used to calculate the least significant difference (LSD). LSD was used regardless of significance of F -ratios in all analyses. Significant differences were sought at the 10% probability level. LSD values at the 5% probability level are included in the tables for readers who prefer to use that level.

The parentage, variety correction factor, and reactions to smut and rust for each clone included in these experiments are listed in table 1. Tables 2 through 6, 7 through 11, and 12 through 16 contain the results of the plant-cane, first-ratoon, and second-ratoon experiments, respectively.

Plant cane

CP 70-1133 yielded significantly more metric tons of cane per hectare (TC/H) than any other clone except CP 79-1380, CP 79-1697, and CP 79-1540 and significantly more metric tons of sugar per hectare (TS/H) than any other clone except CP 79-1697 (tables 2 and 6). CP 79-1697 was lower, but not significantly, than CP 70-1133 in both of these characteristics. Attempts will be made to use CP 79-1697 in future breeding seasons at Canal Point due to its exotic germplasm; it is a BC₁ with parentage of CP 65-357 x US 72-1151 [*Chapina* (a typical *Saccharum officinarum*) x US 56-15-8 (*Saccharum spontaneum*)].

First-Ratoon Cane

CP 78-1247 was the only clone which yielded significantly more TS/H than CP 70-1133 (table 11). The TC/H yield of CP 78-1247 was lower, although not significantly, than that of CP 70-1133 (Table 7); and the yield in kilograms of sugar per metric ton of cane (KS/T) of CP 78-1247 was significantly higher than that of CP 70-1133 (table 10). Results for CP 78-1247 were similar in the plant crop (Glaz et al. 1983) except that its TS/H yield was higher, but not significantly, than that of CP 70-1133, and significantly higher than that of any of the other clones. This year, the TS/H yields of CP 78-1628 and CP 78-2114 were not significantly less than the yield for CP 78-1247 (table 11). Stability of CP 78-1247 for KS/T, TC/H, and TS/H in a

previous selection stage was reported to be high (Kang and Miller 1984). Similar results were reported in plant cane of this selection stage last year (Glaz et al. 1983). However, the results for TS/H in the first-ratoon crop reported herein show that CP 78-1247 was outstanding at three locations (New Farm, Inc., Beardsley, and Lykes Bros.), slightly above average at three locations (Okeelanta, Duda, and South Florida Industries), and slightly less than average at one location (Wedgworth) (table 11), which would indicate that its stability was low (table 11).

CP 78-1628 and CP 78-2114 had higher, although not significantly, yields of TS/H than CP 70-1133. Neither of these new clones had a significantly lower TS/H yield than CP 70-1133 in the plant crop (Glaz et al. 1983). This year, both CP 78-1628 and CP 78-2114 had lower yields of TC/H than CP 70-1133, although only CP 78-1628 was significantly lower (table 7). Also, CP 78-1628 had a significantly higher yield of KS/T than CP 78-2114, which had a significantly higher KS/T yield than CP 70-1133 (table 10).

CP 78-1610, CP 78-1140, CP 78-1156, and CP 78-1263 had TS/H yields not significantly less than the yield for CP 70-1133 (table 11). CP 78-1610 had a TC/H yield not significantly less than that of CP 70-1133 and a KS/T yield significantly higher than that of CP 70-1133 (tables 7 and 10). An interesting characteristic of CP 78-1610 was that it ranked second, and was significantly higher than 10 other clones, in preharvest KS/T but was eighth and significantly higher than only 3 clones in normal harvest KS/T (tables 8 and 10). This relatively high KS/T from early samples caused CP 78-1610 to have the highest estimated TS/H yield from early samples (table 9).

CP 78-1140 had a very high yield of TC/H and a very low yield of KS/T (tables 7 and 10). CP 78-1156 and CP 78-1263 both had moderate yields of TC/H and KS/T (tables 7 and 10).

The data for disease susceptibility indicate that all the above-mentioned clones have adequate resistance for commercial production in Florida. CP 78-1628 currently is classified as intermediate in susceptibility to sugarcane smut (table 1).

Second-Ratoon Cane

CP 77-1414 and CP 77-1055 were the only clones that had significantly more TS/H than CP 63-588 (table 16). Both of these clones also had significantly more TC/H than CP 63-588, and CP 77-1414 had significantly more TC/H than any other clone (table 12). However, CP 77-1414 had significantly less KS/T than CP 63-588 and CP 77-1055, both of which were equal in this characteristic (table 15). In these reports summarizing the results of the past two harvest seasons (Glaz et al. 1982 and 1983), it was reported that CP 77-1414 did not meet the minimal commercial variety correction factor (VCF) requirement of 0.96. This is no longer a valid reason for precluding the approval of CP 77-1414 as a commercial clone. As plant cane, CP 77-1414 had a VCF of 0.928. However, all the samples that were used to calculate that VCF were taken from fields which had previously been frozen. As first- and second-ratoon cane, the VCF's of CP 77-1414 were 0.9969 and 1.0169, respectively, which, including the plant-cane VCF, gave CP 77-1414 an average VCF of 0.9773 (table 1). CP 77-1414 has also had excellent resistance to rust and smut (table 1). CP 77-1055 is susceptible to smut and to a lesser degree, rust (table 1), which detracts from its otherwise

SUMMARY

good production records from plant cane through second-ratoon cane (Glaz et al. 1982 and 1983).

CP 77-1776 has been included in the Florida Sugar Cane League, Inc., seed increase program due to its high yield of KS/T and moderate yields of TC/H as plant cane and first-ratoon cane (Glaz et al. 1982 and 1983). As second-ratoon cane, CP 77-1776 was once again significantly higher than CP 63-588 in early and harvest KS/T (tables 13 and 15). However, the TC/H of CP 77-1776 was significantly less than that of CP 63-588 (table 12), causing CP 77-1776 to have disappointing estimated yields of TS/H from early samples and at harvest time (tables 14 and 16).

CP 77-1776 is resistant to both smut and rust (table 1).

Results from the plant-cane experiments revealed no new high-yielding clones. However, CP 79-1697 was identified as a potentially important clone for breeding due to its nearly acceptable yields and exotic germplasm.

Combined data of the last 2 years (1983-84) indicated seven promising clones from the first-ratoon experiments: CP 78-1140, CP 78-1156, CP 78-1247, CP 78-1263, CP 78-1610, CP 78-1628, and CP 78-2114. Four of these clones, CP 78-1156, CP 78-1247, CP 78-1628, and CP 78-2114, have been included in the Florida Sugar Cane League, Inc., seed increase program. CP 78-1247 has had higher yields of TS/H than CP 70-1133 as plant cane and first-ratoon cane. CP 78-1628 and CP 78-2114 have had TS/H yields about equal to those of CP 70-1133 in the past 2 years. The other three clones yielded significantly less TS/H than CP 70-1133 in plant cane, but not significantly different TS/H in the first-ratoon crop. Of these, CP 78-1610 may be an acceptable early-maturing clone.

After 3 years of evaluation (1982-84), CP 77-1414 had the highest yields of TS/H in the second-ratoon experiments. However, it had lower than average yields of KS/T. CP 77-1776 has had excellent early and harvest yields of KS/T from plant through second-ratoon crop, but its 3-year total of TS/H was only 92.3% of that of CP 63-588. CP 77-1776 has been included in the Florida Sugar Cane League, Inc. seed increase program.

REFERENCES

- Arceneaux, G.
1935. A simplified method of making theoretical sugar yield calculations. In accordance with Winter-Carp-Geerligs formula. *Int. Sugar J.* 37: 264-265.
- Donovan, W.C. and Glaz, B.
1983. Florida's 1983 sugar cane variety census. *Sugar Azucar* 78(12): 42-43, 45-46.
- Glaz, B.; Dean, J. L.; Kang, M. S. and others.
1982. Sugarcane variety tests in Florida. 1981-82 Harvest season. 21 pp. U.S. Agricultural Research Service, New Orleans.
- Glaz, B.; Kang, M. S.; Miller, J. D. and others.
1983. Sugarcane variety tests in Florida. 1982-83 Harvest season. 21 pp. U.S. Agricultural Research Service, New Orleans.
- Kang, M. S. and Miller, J. D.
1984. Genotype x environment interactions for cane and sugar yield and their implications in sugarcane breeding. *Crop Sci.* 24: 435-440.
- Mangelsdorf, A. J.
1983. Cytoplasmic diversity in relation to pests and pathogens. *Sugarcane Breeders' Newsletter* 45: 45-49.
- McIntosh, M. S.
1983. Analysis of combined experiments. *Agron. J.* 75: 153-155.
- Rice, E. R., and Hebert, L. P.
1972. Sugarcane variety tests in Florida during the 1971-72 season. U.S. Agric. Res. Serv. (Rep.) ARS-S-2, 14 pp.
- Snyder, G. H.; Burdine, H. W.; Crockett, J. R. and others.
1978. Water table management for organic soil conservation and crop production in the Florida Everglades. Univ. Fla. Inst. Food Agric. Sci. Tech. Bull. 801, 22 pp.

Table 1.--Parentage, variety correction factors (VCF)¹ and ratings for smut and rust susceptibility of CP 63-588, CP 70-1133, and 32 new sugarcane clones

| Clone | Parentage ² | VCF | Rating ³ | |
|-------------------------------|-----------------------------------|--------|---------------------|------|
| | | | Smut | Rust |
| CP 63-588..... | CL 54-191 X CP 57-120..... | 1.0000 | R | I |
| CP 70-1133..... | ⁴ 67 P 6 CP 56-63..... | .9890 | R | R |
| CP 77-1008..... | CP 65-357 x CL 54-1910..... | .9540 | R | S |
| CP 77-1049..... | CP 68-1154 x CP 68-1022..... | .9980 | I | R |
| CP 77-1055..... | CP 68-1154 x CP 68-1022..... | .9792 | S | I |
| CP 77-1125..... | CP 63-588 x CP 56-63..... | .9700 | R | R |
| CP 77-1148..... | CP 65-357 x CP 68-1022..... | .9610 | I | R |
| CP 77-1400..... | CP 70-1133 x CP 69-1059..... | .9890 | I | R |
| CP 77-1404..... | CP 68-1067 x CP 69-1056..... | .9700 | R | R |
| CP 77-1414..... | CP 68-1067 x CP 69-1056..... | .9773 | R | R |
| CP 77-1446..... | CP 69-1062 x CP 63-306..... | .9870 | S | S |
| CP 77-1720..... | CP 68-1154 x CP 63-588..... | 1.0120 | S | R |
| CP 77-1776 ⁵ | CP 68-1067 x CP 68-1022..... | 1.0302 | R | R |
| CP 78-1038..... | 73 P 1 CP 63-588..... | 1.0205 | I | R |
| CP 78-1140..... | 73 P 2 CP 69-1052..... | .9555 | R | R |
| CP 78-1156 ⁵ | CP 70-1512 x CP 70-1133..... | .9790 | R | R |
| CP 78-1247 ⁵ | CP 68-1067 x CP 57-614..... | .9570 | R | R |
| CP 78-1263..... | CP 68-1067 x CP 57-614..... | .9815 | R | R |
| CP 78-1599..... | CP 68-10k67 x CP 68-1022..... | 1.0025 | I | R |
| CP 78-1610..... | CP 68-1067 x CP 68-1022..... | .9548 | R | R |
| CP 78-1628 ⁵ | CP 65-357 x CP 68-1026..... | .9677 | I | R |
| CP 78-1979..... | CP 70-1512 x CP 68-1026..... | .9759 | I | S |
| CP 78-2114 ⁵ | Unknown..... | 1.0033 | R | R |
| CP 79-1243..... | CP 71-1442 x CP 70-1133..... | .9595 | I | S |
| CP 79-1288..... | CP 73-1372 x CP 70-1133..... | .9783 | R | R |
| CP 79-1374..... | CP 71-1194 x CP 68-1026..... | .9512 | R | R |
| CP 79-1380..... | CP 73-1372 x US 75-1337..... | .9170 | I | R |
| CP 79-1540..... | CP 70-1547 x CP 70-1133..... | .9262 | R | R |
| CP 79-1580..... | CP 73-1225 x CP 72-1370..... | .9408 | R | S |
| CP 79-1606..... | CP 71-1273 x CP 69-1052..... | .9382 | R | R |
| CP 79-1608..... | CP 71-1273 x CP 69-1052..... | .9335 | R | R |
| CP 79-1658..... | CP 70-1547 x CP 63-588..... | 1.0578 | R | R |
| CP 79-1661..... | CP 70-1547 x CP 63-588..... | 1.0022 | R | R |
| CP 79-1697..... | CP 65-357 x US 72-1151..... | .9874 | R | R |

¹VCF used to calculate theoretical yield of 96° sugar per metric ton of cane according to Arceneaux's simplification of the Winter-Carp-Geerligs formula.

²CL 54-191 and CL 54-1910 developed by the U. S. Sugar Corp., Clewiston, Fla.

³R = resistant enough for commercial production; S = too susceptible for commercial production; I = intermediate (the available data not sufficiently persuasive to determine susceptibility).

⁴67 P 6, 6th polycross made in the 1967 crossing season. Female parent (CP 56-63) exposed to pollen from many male varieties; therefore, male parent of CP 70-1133 unknown. Similar explanations for CP 78-1038 and CP 78-1140.

⁵Seed cane currently being increased by Florida Sugar Cane League, Inc., for potential release.

Table 2.--Yields of cane, in metric tons per hectare, from plant cane on Lauderdale, Pahokee, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | Average yield, all farms |
|------------------------------------|--|-----------------------------|--|---------------------|----------------------|------------------------|--------|-----------------------------------|
| | Lauderhill muck | | Pahokee muck | Torry muck | Pompano fine sand | | | |
| | Saunders ¹ 1/17/84 | Okeelanta 2/5/84 | | | | | | |
| | Duda 2/23/84 | New Farm, Inc. 1/7/84 | South Florida Industries 1/22/84 | Wedgworth 2/2/84 | Eastgate 2/27/84 | Lykes Bros. 11/3/84 | | |
| CP 70-1133..... ² 57.93 | 136.85 | 185.35 | 113.12 | 152.46 | 138.76 | 153.38 | 106.86 | 132.93 |
| CP 79-1380..... 52.56 | 130.46 | 168.60 | 114.45 | 131.31 | 134.91 | 153.89 | 115.70 | 125.24 |
| CP 79-1697..... 72.98 | 124.46 | 158.24 | 124.63 | 128.36 | 136.60 | 139.15 | 113.19 | 124.70 |
| CP 79-1540..... 71.09 | 117.59 | 148.62 | 120.38 | 147.23 | 119.20 | 149.71 | 120.65 | 124.31 |
| CP 79-1608..... ² 42.99 | 117.26 | 154.55 | 120.02 | 133.22 | 138.92 | 136.62 | 86.01 | 116.20 |
| CP 79-1580..... ² 23.49 | 128.29 | 143.83 | 110.65 | 130.10 | 130.16 | 134.24 | 98.42 | 115.27 |
| CP 79-1606..... 38.12 | 121.45 | 148.10 | 103.41 | 125.13 | 123.65 | 120.67 | 92.48 | 109.13 |
| CP 79-1661..... ² 62.81 | 85.55 | 131.45 | 105.93 | 107.32 | 101.63 | 151.93 | 108.63 | 108.33 |
| CP 79-1288..... 57.65 | 104.89 | 130.42 | 88.78 | 115.55 | 125.27 | 124.84 | 109.99 | 107.17 |
| CP 79-1658..... 84.58 | 91.73 | 133.01 | 89.39 | 107.11 | 81.28 | 120.70 | 106.52 | 101.79 |
| CP 79-1374..... 73.33 | 77.25 | 115.38 | 85.73 | 105.56 | 73.47 | 148.10 | 110.82 | 98.71 |
| CP 79-1243..... 41.88 | 93.77 | 123.32 | 93.59 | 91.28 | 98.88 | 133.42 | 101.18 | 97.16 |
| Mean ³ 57.19 | 110.80 | 145.07 | 105.84 | 122.89 | 116.89 | 138.89 | 105.87 | 113.37 |
| LSD: | | | | | | | | |
| 5% level .. 30.34 | 10.01 | 21.80 | 12.72 | 16.01 | 22.18 | 14.97 | 12.04 | 13.53 |
| 10% level.. 25.22 | 8.32 | 18.12 | 10.57 | 13.31 | 18.43 | 12.44 | 10.00 | 11.29 |
| CV ⁴ (%)..... 36.75 | 6.26 | 10.41 | 8.32 | 9.02 | 13.14 | 7.46 | 7.87 | 11.27 |

¹Flooded conditions after planting reduced emergence considerably at Saunders.

²Average of 3 not 4 replications.

³LSD for location means=5.58 t/ha at 10% probability level.

⁴CV=coefficient of variation.

Table 3.--Indicated yields of 96° sugar, in kilograms per metric ton of cane, from preharvest samples of plant cane on Lauderdalehill, Pahokee, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and sampling date ¹ | | | | | | Average yield, all farms | |
|---------------------------|--|-----------------------|------------------|---|-----------------------|-------------------------|-----------------------------------|-------|
| | Lauderhill muck | | Pahokee muck | | Torry muck | Pompano fine sand | | |
| | Saunders 10/19/83 | Okeelanta 10/20/83 | Duda 10/26/83 | South Florida Industries 10/24/83 | Wedgworth 10/26/83 | Lykes Bros. 10/19/83 | | |
| CP 79-1658..... | 103.7 | 104.5 | 82.5 | 113.5 | 112.0 | 81.4 | 101.0 | 99.8 |
| CP 79-1243..... | 100.0 | 98.5 | 91.6 | 100.7 | 110.0 | 66.8 | 116.8 | 97.8 |
| CP 70-1133..... | 105.6 | 80.7 | 84.7 | 104.8 | 92.4 | 101.1 | 114.6 | 97.7 |
| CP 79-1288..... | 105.3 | 92.3 | 78.3 | 110.2 | 99.7 | 74.3 | 113.1 | 96.2 |
| CP 79-1697..... | 97.8 | 73.7 | 69.7 | 110.9 | 90.5 | 80.4 | 98.4 | 88.8 |
| CP 79-1580..... | 91.1 | 76.8 | 88.5 | 87.8 | 90.6 | 69.3 | 101.3 | 86.5 |
| CP 79-1606..... | 99.7 | 89.5 | 73.2 | 80.3 | 90.5 | 68.4 | 101.0 | 86.1 |
| CP 79-1380..... | 86.5 | 85.4 | 76.3 | 87.7 | 84.8 | 86.8 | 94.3 | 85.9 |
| CP 79-1661..... | 91.6 | 84.5 | 75.6 | 80.4 | 83.0 | 71.0 | 97.4 | 83.4 |
| CP 79-1374..... | 93.4 | 80.1 | 69.6 | 90.6 | 78.1 | 62.8 | 98.1 | 81.8 |
| CP 79-1608..... | 86.7 | 84.2 | 68.3 | 80.9 | 89.1 | 64.9 | 91.4 | 80.8 |
| CP 79-1540..... | 80.5 | 62.6 | 67.8 | 95.9 | 90.5 | 74.0 | 88.3 | 79.9 |
| Mean ¹ | 95.2 | 84.4 | 77.2 | 95.3 | 92.6 | 74.6 | 101.3 | 88.7 |
| LSD: | | | | | | | | |
| 5% level..... | 17.6 | 25.3 | 18.0 | 38.7 | 54.9 | 47.1 | 10.4 | 6.2 |
| 10% level..... | 14.3 | 20.6 | 14.7 | 31.5 | 44.8 | 38.5 | 8.5 | 5.1 |
| CV ² (%) | 8.38 | 13.62 | 10.59 | 18.43 | 26.95 | 28.71 | 4.66 | 14.53 |

¹LSD for location means=5.42 kg/t of cane at 10% probability level.

²CV=coefficient of variation.

Table 4.--Indicated yields of 96° sugar, in metric tons per hectare, from preharvest samples of plant cane on
Lauderhill, Pahokee, and Torry muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and sampling date ¹ | | | | | | Average yield, all farms |
|--------------------------|--|-----------------------|------------------|---|-----------------------|----------------------|-----------------------------------|
| | Lauderhill muck | | Pahokee muck | | Torry muck | Pompano fine sand | |
| | Saunders ² 10/19/83 | Okeelanta 10/20/83 | Duda 10/26/83 | South Florida Industries 10/24/83 | Wedgworth 10/26/83 | Eastgate 10/17/83 | Lykes Bros. 10/19/83 |
| CP 70-1133..... | 3.894 | 11.081 | 14.648 | 16.886 | 11.074 | 16.220 | 12.050 |
| CP 79-1697..... | 5.590 | 9.258 | 11.878 | 14.598 | 12.629 | 11.916 | 10.326 |
| CP 79-1380..... | 4.174 | 11.696 | 13.213 | 11.908 | 10.845 | 13.547 | 11.162 |
| CP 79-1288..... | 6.990 | 9.489 | 9.899 | 12.649 | 11.771 | 8.910 | 11.858 |
| CP 79-1658..... | 8.366 | 9.973 | 11.390 | 11.840 | 9.618 | 9.689 | 10.338 |
| CP 79-1540..... | 4.784 | 7.600 | 10.225 | 14.003 | 12.018 | 10.051 | 11.286 |
| CP 79-1580..... | 1.889 | 9.841 | 12.314 | 11.475 | 12.265 | 9.140 | 9.632 |
| CP 79-1243..... | 3.724 | 9.711 | 11.291 | 9.208 | 11.928 | 8.828 | 11.828 |
| CP 79-1608..... | 3.764 | 10.017 | 10.278 | 11.557 | 12.616 | 8.472 | 7.353 |
| CP 79-1606..... | 4.488 | 10.911 | 10.321 | 9.157 | 10.881 | 7.554 | 9.389 |
| CP 79-1661..... | 4.853 | 7.904 | 10.009 | 8.196 | 8.726 | 10.806 | 10.287 |
| CP 79-1374..... | 7.274 | 6.513 | 8.475 | 9.389 | 6.083 | 9.247 | 10.803 |
| Mean ³ | 4.983 | 9.499 | 11.162 | 11.739 | 10.871 | 10.227 | 10.526 |
| LSD: | | | | | | | |
| 5% level..... | 5.495 | 3.383 | 3.034 | 4.691 | 3.676 | 7.225 | 1.510 |
| 10% level..... | 4.484 | 2.761 | 2.476 | 3.828 | 2.999 | 5.895 | 1.232 |
| CV ⁴ (%)..... | 50.11 | 16.18 | 12.35 | 18.15 | 15.36 | 32.10 | 6.52 |
| | | | | | | | 20.46 |
| | | | | | | | 9.856 |
| | | | | | | | 1.818 |
| | | | | | | | 1.519 |
| | | | | | | | 8.957 |
| | | | | | | | 8.683 |
| | | | | | | | 8.255 |

¹Yields based on early sucrose analysis, assuming that early cane yields are equal to actual yields at harvest.

²Flooded conditions after planting reduced emergence considerably at Saunders.

³LSD for location means=0.956 t/ha at 10% probability level.

⁴CV=coefficient of variation.

Table 5.--Indicated yields of 96° sugar, in kilograms per metric ton of cane, from plant cane on Lauderhill, Pahokee, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | Average yield, all farms | |
|--------------------------|--|---------------------|-----------------|-----------------------------|--|----------------------|---------------------|-----------------------------------|-----------------------|
| | Lauderhill muck | | Pahokee muck | | Torry muck | Pompano fine sand | | | |
| | Saunders 1/17/84 | Okeelanta 2/5/84 | Duda 2/23/84 | New Farm, Inc. 1/7/84 | South Florida Industries 1/22/84 | Wedgworth 2/2/84 | Eastgate 2/27/84 | | Lykes Bros. 1/3/84 |
| CP 79-1243..... | 128.8 | 126.9 | 114.9 | 119.3 | 118.5 | 116.2 | 124.5 | 140.5 | 123.7 |
| CP 79-1658..... | 120.9 | 108.0 | 94.8 | 123.6 | 127.3 | 99.1 | 123.5 | 141.4 | 117.3 |
| CP 79-1288..... | 125.3 | 106.0 | 99.7 | 107.2 | 121.0 | 102.4 | 113.5 | 139.0 | 114.2 |
| CP 79-1606..... | 107.2 | 109.8 | 103.0 | 114.4 | 118.2 | 98.4 | 124.8 | 134.0 | 113.7 |
| CP 70-1133..... | 110.0 | 102.0 | 101.0 | 101.1 | 111.5 | 92.1 | 115.6 | 130.5 | 107.9 |
| CP 79-1580..... | 105.8 | 99.3 | 101.6 | 104.9 | 107.8 | 96.9 | 108.6 | 123.3 | 106.0 |
| CP 79-1697..... | 107.0 | 94.4 | 89.2 | 98.4 | 104.1 | 100.3 | 125.8 | 126.7 | 105.7 |
| CP 79-1661..... | 108.2 | 95.7 | 96.5 | 97.3 | 104.4 | 101.1 | 116.8 | 124.4 | 105.5 |
| CP 79-1374..... | 101.6 | 101.7 | 90.8 | 101.3 | 110.4 | 101.0 | 97.5 | 124.4 | 103.6 |
| CP 79-1380..... | 103.2 | 96.1 | 84.9 | 101.5 | 109.7 | 94.5 | 103.0 | 121.3 | 101.8 |
| CP 79-1608..... | 96.0 | 98.3 | 98.3 | 94.1 | 102.2 | 95.5 | 111.1 | 116.8 | 101.5 |
| CP 79-1540..... | 100.5 | 97.2 | 93.6 | 85.4 | 102.0 | 94.1 | 97.2 | 117.8 | 98.5 |
| Mean ¹ | 109.6 | 102.9 | 97.4 | 104.0 | 111.4 | 99.3 | 113.5 | 128.3 | 108.3 |
| LSD: | | | | | | | | | |
| 5% level..... | 8.1 | 7.8 | 11.8 | 10.3 | 12.2 | 10.1 | 12.7 | 5.5 | 5.4 |
| 10% level.... | 6.7 | 6.5 | 9.8 | 8.6 | 10.1 | 8.4 | 10.5 | 4.6 | 4.5 |
| CV ² (%)..... | 5.10 | 5.26 | 8.42 | 6.88 | 7.57 | 7.07 | 7.73 | 2.97 | 6.47 |

¹LSD for location means=2.5 kg/t of cane at 10% probability level.

²CV=coefficient of variation.

Table 6.--Indicated yields of 96° sugar, in metric tons per hectare, from plant cane on Lauderdale, Pahokee, and Torry muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | | Average yield, all farms |
|------------------------------------|--|---------------------|-----------------|-----------------------------|--|---------------------|--------------------------|---|-----------------------------------|
| | Lauderhill muck | | | | Pahokee muck | | | | |
| | Saunders ¹ 1/17/84 | Okeelanta 2/5/84 | Duda 2/23/84 | New Farm, Inc. 1/7/84 | South Florida Industries 1/22/84 | Wedgworth 2/2/84 | Torry muck 2/27/84 | Pompano fine sand Lykes Bros. 1/3/84 | |
| CP 70-1133..... ² 6.302 | 13.950 | 18.720 | 11.425 | 16.974 | 12.759 | 17.750 | 13.937 | 14.225 | |
| CP 79-1697..... 7.807 | 11.776 | 14.186 | 12.324 | 13.413 | 13.700 | 17.552 | 14.319 | 13.134 | |
| CP 79-1380..... 5.441 | 12.522 | 14.287 | 11.634 | 14.390 | 12.788 | 15.851 | 14.019 | 12.616 | |
| CP 79-1606..... 4.025 | 13.343 | 15.268 | 11.841 | 14.835 | 12.180 | 15.097 | 12.403 | 12.374 | |
| CP 79-1540..... 7.114 | 11.429 | 14.064 | 10.313 | 14.995 | 11.147 | 14.534 | 14.227 | 12.228 | |
| CP 79-1580..... ² 2.451 | 12.726 | 14.608 | 11.601 | 14.010 | 12.604 | 14.583 | 12.139 | 12.143 | |
| CP 79-1288..... 7.210 | 11.125 | 12.933 | 9.513 | 13.990 | 12.881 | 14.148 | 15.277 | 12.135 | |
| CP 79-1243..... 5.431 | 11.891 | 14.148 | 11.168 | 10.841 | 11.488 | 16.685 | 14.204 | 11.982 | |
| CP 79-1658..... ¹⁰ 2.38 | 9.949 | 12.589 | 11.094 | 13.644 | 8.040 | 14.896 | 15.052 | 11.938 | |
| CP 79-1608..... 4.181 | 11.516 | 15.180 | 11.258 | 13.702 | 13.244 | 15.174 | 10.030 | 11.786 | |
| CP 79-1661..... ² 6.730 | 8.132 | 12.655 | 10.295 | 11.159 | 10.259 | 17.753 | 13.516 | 11.460 | |
| CP 79-1374..... 7.392 | 7.866 | 10.457 | 8.647 | 11.682 | 7.403 | 14.446 | 13.784 | 10.210 | |
| Mean ³ 6.262 | 11.352 | 14.091 | 10.926 | 13.636 | 11.541 | 15.706 | 13.576 | 12.183 | |
| LSD: | | | | | | | | | |
| 5% level.... 3.203 | 1.455 | 2.703 | 1.818 | 2.461 | 2.464 | 2.665 | 1.540 | 1.530 | |
| 10% level... 2.662 | 1.209 | 2.246 | 1.511 | 2.045 | 2.047 | 2.215 | 1.280 | 1.278 | |
| CV ⁴ (%).....35.43 | 8.87 | 13.28 | 11.52 | 12.50 | 14.78 | 11.75 | 7.855 | 13.36 | |

¹Flooded conditions after planting reduced emergence considerably at Saunders.

²Average is of 3 not 4 replications.

³LSD for location means=0.598 t/ha at 10% probability level.

⁴CV=coefficient of variation.

Table 7.--Yields of cane, in metric tons per hectare, from first-ratoon cane on Lauderhill, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | Average yield, all farms | |
|--------------------------|--|----------------------|-----------------|--|----------------------|-----------------------------|----------------------|-----------------------------------|-----------------------|
| | Lauderhill muck | | Pahokee muck | Terra Ceia muck | Torry muck | Pompano fine sand | | | |
| | Saunders 1/17/84 | Okeelanta 2/16/84 | Duda 2/24/84 | South Florida Industries 1/21/84 | Wedgworth 12/7/83 | New Farm, Inc. 1/1/84 | Beardsley 2/11/84 | | Lykes Bros. 1/4/84 |
| CP 78-1140..... | 109.19 | 98.15 | 142.61 | 137.84 | 111.79 | 114.11 | 147.39 | 98.99 | 120.01 |
| CP 70-1133..... | 112.01 | 100.16 | 141.02 | 129.92 | 105.68 | 123.83 | 152.75 | 89.98 | 119.42 |
| CP 78-2114..... | | 103.09 | 127.73 | 117.74 | 109.73 | 120.90 | 137.52 | 98.36 | 116.44 |
| CP 78-1247..... | | 92.78 | 118.10 | 118.06 | 98.20 | 123.65 | 152.52 | 108.54 | 115.98 |
| CP 78-1610..... | | 104.28 | 123.46 | 132.05 | 97.29 | 109.94 | 150.12 | 90.86 | 115.43 |
| CP 78-1628..... | 106.48 | 97.81 | 123.28 | 125.74 | 98.68 | 118.14 | 135.15 | 96.14 | 112.68 |
| CP 78-1263..... | 98.19 | 102.81 | 117.15 | 130.36 | 105.14 | 114.39 | 126.65 | 88.80 | 110.44 |
| CP 78-1156..... | | 94.21 | 112.72 | 116.94 | 108.74 | 99.75 | 149.05 | 90.50 | 110.27 |
| CP 78-1979..... | | 86.75 | 105.12 | 109.08 | 89.65 | 105.18 | 113.44 | 88.65 | 98.70 |
| CP 78-1599..... | | | 101.27 | 111.14 | 99.52 | 93.46 | 105.68 | 73.02 | 97.35 |
| CP 63-588..... | 73.59 | 79.17 | 102.17 | 114.59 | 98.55 | 88.88 | 115.80 | 87.24 | 95.00 |
| CP 78-1038..... | | 89.41 | 111.62 | 104.40 | 89.25 | 88.14 | 129.81 | 45.89 | 94.07 |
| Mean ¹ | 99.89 | 95.33 | 118.85 | 120.65 | 101.02 | 108.36 | 134.66 | 88.08 | 109.10 |
| LSD: | | | | | | | | | |
| 5% level.... | 12.28 | 9.95 | 9.56 | 14.90 | 23.61 | 14.81 | 16.18 | 19.30 | 5.41 |
| 10% level... | 10.05 | 8.27 | 7.94 | 12.38 | 19.62 | 12.31 | 13.44 | 16.04 | 4.54 |
| CV ² (%)..... | 7.98 | 7.07 | 5.57 | 8.57 | 16.18 | 9.47 | 8.29 | 15.18 | 10.11 |

¹LSD for location means=3.71 t/ha at 10% probability level.

²CV=coefficient of variation.

Table 8.--Indicated yields of 96° sugar, in kilograms per metric ton of cane, from preharvest samples of first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and sampling date | | | | | | Average yield, all farms | |
|--------------------------|---|--------------------------|-----------------|-------------|-------------------|-------------|--------------------------|-------|
| | Lauderhill muck | Pahokee muck | Terra Ceia muck | Torrey muck | Pompano fine sand | | | |
| | Saunders Okeelanta Duda | South Florida Industries | New Farm, Inc. | Wedgworth | Beardsley | Lykes Bros. | | |
| | 10/19/83 10/20/83 10/26/83 | 10/24/83 | 10/24/83 | 10/26/83 | 10/17/83 | 10/19/83 | | |
| CP 78-1599..... | | 120.1 | 110.3 | 116.2 | 118.6 | 108.5 | 124.6 | 116.4 |
| CP 78-1610..... | 97.3 | 102.7 | 110.0 | 112.7 | 114.2 | 96.5 | 116.8 | 107.2 |
| CP 78-1247..... | 76.9 | 123.4 | 106.3 | 103.2 | 101.9 | 84.8 | 116.4 | 101.8 |
| CP 78-1156..... | 91.7 | 91.5 | 108.7 | 98.2 | 103.3 | 97.4 | 108.9 | 100.0 |
| CP 78-1038..... | 95.9 | 108.4 | 110.0 | 100.2 | 101.8 | 58.8 | 122.0 | 99.6 |
| CP 78-2114..... | 92.6 | 87.5 | 83.8 | 93.3 | 98.0 | 98.9 | 123.6 | 96.8 |
| CP 70-1133..... | 88.9 | 84.0 | 97.8 | 85.6 | 98.3 | 97.4 | 125.0 | 96.3 |
| CP 78-1263..... | 93.2 | 94.6 | 96.8 | 93.8 | 94.5 | 73.5 | 110.5 | 92.7 |
| CP 78-1140..... | 89.0 | 101.7 | 86.2 | 95.6 | 87.2 | 93.4 | 99.4 | 91.7 |
| CP 78-1628..... | 85.2 | 83.9 | 96.9 | 96.8 | 89.3 | 78.3 | 109.6 | 91.1 |
| CP 63-588..... | 83.4 | 97.0 | 86.5 | 87.3 | 90.1 | 71.9 | 107.5 | 89.7 |
| CP 78-1979..... | | 78.6 | 83.0 | 81.6 | 95.9 | 70.4 | 86.7 | 82.5 |
| Mean ¹ | 87.9 | 97.8 | 98.9 | 97.1 | 99.4 | 85.8 | 112.6 | 96.7 |
| LSD: | | | | | | | | |
| 5% level... | 29.7 | 25.3 | 22.1 | 15.4 | 21.0 | 21.0 | 18.2 | 7.0 |
| 10% level... | 22.8 | 20.7 | 18.0 | 12.6 | 17.1 | 17.2 | 14.9 | 5.8 |
| CV ² (%)..... | 12.2 | 11.77 | 10.17 | 7.21 | 9.60 | 11.14 | 7.36 | 10.28 |

¹LSD for location means=5.7 kg/t of cane at 10% probability level.

²CV=coefficient of variation.

Table 9.--Indicated yields of 96° sugar, in metric tons per hectare, from preharvest samples of first-ratoon cane on Lauderdale, Pahokee, Terra Ceia, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and sampling date ¹ | | | | | | Average yield, all farms | |
|--------------------------|--|--------------------|---------------|-----------------------------------|-------------------------|--------------------|--------------------------|----------------------|
| | Lauderhill muck | | Pahokee muck | Terra Ceia muck | Torry muck | Pompano fine sand | | |
| | Saunders 10/19/83 | Okeelanta 10/20/83 | Duda 10/26/83 | South Florida Industries 10/24/83 | New Farm, Inc. 10/24/83 | Wedgworth 10/26/83 | Beardsley 10/17/83 | Lykes Bros. 10/19/83 |
| CP 78-1610..... | | 9.970 | 12.626 | 15.101 | 12.097 | 11.198 | 14.498 | 10.889 |
| CP 78-1247..... | | 7.497 | 15.025 | 12.691 | 11.568 | 10.516 | 13.351 | 12.782 |
| CP 70-1133..... | 10.200 | 9.563 | 11.538 | 13.509 | 10.582 | 10.039 | 15.911 | 9.767 |
| CP 78-1140..... | 9.511 | 8.076 | 15.455 | 12.481 | 10.272 | 9.207 | 13.863 | 9.966 |
| CP 78-1599..... | | | 11.657 | 11.242 | 10.200 | 11.759 | 12.435 | 8.518 |
| CP 78-2114..... | | 9.103 | 11.008 | 10.118 | 10.652 | 10.660 | 13.358 | 11.531 |
| CP 78-1156..... | | 8.661 | 10.282 | 12.479 | 9.663 | 10.190 | 14.082 | 9.191 |
| CP 78-1263..... | 9.558 | 7.969 | 11.164 | 12.465 | 11.490 | 12.255 | 9.092 | 10.628 |
| CP 78-1628..... | 9.470 | 8.240 | 10.603 | 12.410 | 11.070 | 8.286 | 10.248 | 9.705 |
| CP 78-1038..... | | 8.391 | 12.258 | 11.083 | 8.984 | 9.188 | 7.607 | 3.349 |
| CP 63-588..... | 6.340 | 7.138 | 9.984 | 10.336 | 7.493 | 7.537 | 7.594 | 8.074 |
| CP 78-1979..... | | 6.926 | 8.032 | 9.107 | 8.000 | 8.521 | 7.316 | 8.025 |
| Mean ² | 9.016 | 8.517 | 11.636 | 11.973 | 10.172 | 9.946 | 11.613 | 9.369 |
| LSD: | | | | | | | | |
| 5% level... | 4.982 | 3.470 | 3.667 | 4.197 | 2.210 | 3.315 | 3.065 | 2.833 |
| 10% level.. | 3.826 | 2.832 | 2.992 | 3.425 | 1.803 | 2.706 | 2.501 | 2.311 |
| CV ³ (%)..... | 19.91 | 18.51 | 14.32 | 15.93 | 9.87 | 15.14 | 11.99 | 13.74 |
| | | | | | | | | 14.60 |
| | | | | | | | | 10.389 |
| | | | | | | | | 0.871 |
| | | | | | | | | 0.727 |
| | | | | | | | | 14.60 |

¹Yields based on early sucrose analysis, assuming that early cane yields are equal to actual yields at harvest.
²LSD for location means=1.628 t/ha at 10% probability level.
³CV=coefficient of variation.

Table 10.--Indicated yields of 96° sugar, in kilograms per metric ton of cane, from first-ratoon cane on
Lauderhill, Pahokee, Terra Ceia, and Torry muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | Average yield, all farms |
|--------------------------|--|---------------|-----------------|---------------|----------------------|-------------|-----------------------------------|
| | Lauderhill muck | Pahokee muck | Terra Ceia muck | Torry muck | Pompano fine sand | | |
| Saunders Okeelanta Duda | | South Florida | Wedgworth | New Farm, | Beardsley | Lykes Bros. | |
| 1/17/84 2/16/84 | 2/24/84 | Industries | 12/7/83 | Inc. | 2/11/84 | 1/4/84 | |
| | | 1/21/84 | | 1/1/84 | | | |
| CP 78-1038..... | 136.5 | 125.5 | 132.1 | 107.2 | 123.3 | 119.8 | 126.3 |
| CP 78-1599..... | | 117.2 | 128.8 | 117.8 | 119.9 | 115.4 | 123.7 |
| CP 78-1247..... | 117.6 | 114.5 | 116.5 | 111.3 | 120.6 | 117.6 | 118.9 |
| CP 63-588..... | 119.8 | 117.9 | 113.9 | 107.7 | 116.9 | 111.1 | 117.7 |
| CP 78-1628..... | 115.6 | 112.7 | 113.4 | 107.4 | 119.8 | 115.8 | 117.5 |
| CP 78-1156..... | 122.0 | 112.0 | 116.5 | 111.2 | 94.2 | 112.9 | 115.6 |
| CP 78-2114..... | 115.1 | 112.0 | 109.6 | 110.0 | 109.8 | 105.0 | 114.1 |
| CP 78-1610..... | 107.2 | 104.1 | 112.8 | 108.7 | 117.3 | 111.3 | 113.8 |
| CP 78-1263..... | 117.4 | 110.8 | 114.1 | 116.1 | 108.3 | 98.6 | 113.6 |
| CP 70-1133..... | 104.9 | 107.2 | 104.1 | 109.2 | 111.3 | 108.0 | 110.5 |
| CP 78-1140..... | 105.5 | 102.8 | 104.7 | 105.3 | 106.9 | 108.6 | 108.6 |
| CP 78-1979..... | 96.9 | 104.5 | 102.9 | 98.7 | 98.7 | 92.5 | 102.5 |
| Mean ¹ | 114.4 | 111.8 | 114.1 | 109.2 | 112.2 | 109.7 | 115.04 |
| LSD: | | | | | | | |
| 5% level.... | 8.8 | 11.6 | 4.3 | 9.3 | 14.2 | 10.6 | 3.3 |
| 10% level.... | 7.2 | 9.7 | 3.6 | 7.8 | 11.8 | 8.9 | 2.8 |
| CV ² (%)..... | 5.18 | 7.22 | 2.66 | 5.93 | 8.78 | 6.73 | 5.89 |

¹LSD for location means=2.3 kg/t of cane at 10% probability level.
²CV=coefficient of variation.

Table 12.--Yields of cane, in metric tons per hectare, from second-ratoon cane on Lauderdale, Terra Ceia, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | Average yield, all farms |
|--------------------------|--|------------------|----------------------|---------------------|-------------------------------------|----------------------|-----------------------|--------------------------|
| | Lauderdale muck | | | Terra Ceia muck | | Torrey muck | Pompano fine sand | |
| | New Farm, Inc. 10/30/83 | Duda 11/14/83 | Wedgworth 12/7/83 | Saunders 1/18/84 | South Florida Industries 1/20/84 | Beardsley 2/12/84 | Lykes Bros. 1/2/84 | |
| CP 77-1414..... | 87.62 | 122.85 | 123.46 | 87.59 | 133.85 | 131.08 | 104.73 | 113.03 |
| CP 77-1055..... | 66.11 | 99.49 | 104.61 | 61.35 | 114.20 | 150.01 | 97.25 | 99.00 |
| CP 77-1008..... | 59.73 | 83.70 | 93.33 | 81.52 | 101.38 | 126.20 | 88.85 | 90.67 |
| CP 77-1125..... | 68.91 | 95.56 | 85.68 | 76.87 | 115.48 | 136.16 | 36.52 | 87.88 |
| CP 77-1404..... | 62.85 | 87.06 | 73.70 | 57.45 | 93.14 | 157.34 | 73.61 | 86.45 |
| CP 77-1148..... | 51.86 | 96.92 | 100.12 | 61.13 | 110.70 | 89.96 | 71.44 | 83.16 |
| CP 63-588..... | 39.26 | 69.05 | 64.30 | 59.77 | 95.78 | 158.45 | 85.91 | 81.79 |
| CP 77-1049..... | 60.51 | 64.82 | 80.37 | 76.99 | 73.16 | 120.80 | 64.61 | 77.32 |
| CP 77-1720..... | 49.19 | 56.96 | 62.22 | 62.85 | 102.63 | 78.98 | 99.59 | 73.20 |
| CP 77-1446..... | 46.19 | 87.31 | 92.58 | 59.78 | 85.46 | 51.95 | 86.16 | 72.78 |
| CP 77-1776..... | 51.20 | 70.83 | 74.60 | 50.15 | 172.56 | 94.59 | 70.96 | 69.27 |
| CP 77-1400..... | 33.99 | 67.55 | 80.27 | 48.00 | 80.04 | 98.09 | 63.82 | 67.40 |
| Mean ¹ | 56.45 | 83.51 | 86.27 | 65.28 | 98.20 | 116.13 | 78.62 | 83.49 |
| LSD: | | | | | | | | |
| 5% level..... | 23.40 | 13.09 | 16.38 | 20.48 | 21.04 | 21.83 | 16.13 | 6.15 |
| 10% level..... | 19.45 | 10.88 | 13.61 | 17.02 | 17.49 | 17.86 | 13.40 | 5.14 |
| CV ² (%)..... | 28.71 | 10.85 | 13.15 | 21.73 | 14.84 | 12.93 | 14.20 | 15.93 |

¹LSD for location means=7.27 t/ha at 10% probability level.
²CV=coefficient of variation.

Table 13.--Indicated yields of 96° sugar, in kilograms per metric ton of cane, from preharvest samples of second-ratoon cane on Lauderhill, Terra Ceia, and Torry muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and sampling date | | | | | | Average yield, all farms | |
|---------------------------|---|-------------------------------|------------------|-----------------------|---|-----------------------|--------------------------|-------------------------|
| | Lauderhill muck | | | Terra Ceia muck | Torry muck | Pompano fine sand | | |
| | Saunders 10/19/83 | New Farm, Inc. 10/24/83 | Duda 10/26/83 | Wedgworth 10/26/83 | South Florida Industries 10/24/83 | Beardsley 10/17/83 | | Lykes Bros. 10/19/83 |
| CP 77-1776..... | 95.4 | 112.3 | 107.7 | 121.8 | 120.5 | 86.9 | 138.1 | 111.8 |
| CP 77-1049..... | 111.9 | 99.2 | 115.5 | 102.0 | 123.2 | 87.4 | 121.6 | 108.7 |
| CP 77-1148..... | 99.7 | 100.8 | 109.6 | 94.2 | 122.9 | 91.7 | 132.7 | 107.4 |
| CP 77-1055..... | 90.1 | 92.7 | 120.0 | 107.8 | 105.1 | 87.8 | 125.7 | 104.2 |
| CP 77-1400..... | 105.8 | 111.1 | 99.4 | 107.8 | 99.2 | 90.7 | 112.2 | 103.7 |
| CP 77-1404..... | 94.6 | 95.1 | 121.4 | 102.3 | 98.1 | 98.1 | 111.1 | 103.0 |
| CP 77-1446..... | 100.1 | 112.1 | 106.2 | 86.4 | 112.6 | 94.8 | 108.5 | 102.9 |
| CP 77-1720..... | 99.3 | 91.3 | 105.1 | 103.9 | 108.1 | 78.1 | 105.0 | 98.7 |
| CP 77-1125..... | 101.7 | 104.0 | 93.8 | 110.5 | 90.3 | 78.4 | 122.2 | 100.1 |
| CP 63-588 | 88.5 | 94.5 | 88.6 | 100.0 | 104.0 | 90.8 | 126.2 | 98.9 |
| CP 77-1414..... | 89.9 | 98.5 | 103.7 | 108.1 | 100.3 | 73.9 | 108.4 | 97.5 |
| CP 77-1008..... | 68.1 | 91.0 | 94.5 | 101.3 | 96.2 | 86.8 | 104.0 | 91.7 |
| Mean ¹ | 95.4 | 100.2 | 105.5 | 103.8 | 106.7 | 87.1 | 118.0 | 102.4 |
| LSD: | | | | | | | | |
| 5% level..... | 20.5 | 28.9 | 33.2 | 30.5 | 13.5 | 25.9 | 28.2 | 9.4 |
| 10% level..... | 16.8 | 23.6 | 29.9 | 24.9 | 11.0 | 21.2 | 23.0 | 7.8 |
| CV ² (%) | 9.78 | 8.69 | 15.76 | 13.36 | 5.73 | 13.47 | 10.85 | 12.06 |

¹LSD for location means=10.75 kg/t of cane at 10% probability level.

²CV=coefficient of variation.

Table 14.--Indicated yields of 96° sugar, in metric tons per hectare, from preharvest samples of second-ratoon cane on Lauderdale, Terra Ceia, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and sampling date ¹ | | | | | | Average yield, all farms | |
|--------------------------|--|-------------------------------|--------------------|-----------------------|---|-----------------------|-----------------------------------|-------------------------|
| | Lauderhill muck | | Terra Ceia muck | Torry muck | Pompano fine sand | | | |
| | Saunders 10/19/83 | New Farm, Inc. 10/24/83 | Duda 10/26/83 | Wedgworth 10/26/83 | South Florida Industries 10/24/83 | Beardsley 10/17/83 | | Lykes Bros. 10/19/83 |
| CP 77-1414..... | 7.700 | 10.821 | 12.849 | 13.430 | 13.135 | 9.778 | 11.145 | 11.265 |
| CP 77-1055..... | 6.747 | 5.576 | 12.392 | 11.510 | 10.983 | 12.797 | 11.449 | 10.207 |
| CP 77-1148..... | 6.365 | 4.363 | 10.863 | 8.998 | 11.900 | 8.909 | 10.231 | 8.804 |
| CP 77-1404..... | 5.823 | 6.894 | 10.040 | 6.923 | 8.889 | 14.668 | 6.555 | 8.542 |
| CP 77-1125..... | 8.270 | 7.990 | 8.875 | 9.171 | 10.139 | 11.115 | 4.228 | 8.541 |
| CP 77-1049..... | 9.441 | 6.791 | 8.902 | 7.742 | 9.351 | 10.176 | 6.541 | 8.420 |
| CP 77-1008..... | 4.801 | 6.899 | 7.939 | 9.342 | 10.046 | 10.259 | 8.573 | 8.266 |
| CP 63-588..... | 5.218 | 3.587 | 6.411 | 6.423 | 10.131 | 14.202 | 9.945 | 7.988 |
| CP 77-1776..... | 4.522 | 6.040 | 8.166 | 8.697 | 7.154 | 8.835 | 9.071 | 7.498 |
| CP 77-1446..... | 5.843 | 5.271 | 8.509 | 7.798 | 9.559 | 5.782 | 8.859 | 7.375 |
| CP 77-1720..... | 6.960 | 5.927 | 6.085 | 5.285 | 10.820 | 3.774 | 10.238 | 7.013 |
| CP 77-1400..... | 4.655 | 3.594 | 7.089 | 7.957 | 7.178 | 9.561 | 7.843 | 6.840 |
| Mean ² | 6.362 | 6.146 | 9.010 | 8.606 | 9.941 | 9.988 | 8.723 | 8.396 |
| LSD: | | | | | | | | |
| 5% level..... | 2.757 | 3.267 | 3.225 | 4.243 | 3.443 | 4.105 | 3.483 | 1.971 |
| 10% level..... | 2.093 | 2.666 | 2.632 | 3.462 | 2.809 | 3.539 | 2.842 | 1.647 |
| CV ³ (%)..... | 19.69 | 24.15 | 16.26 | 22.40 | 15.73 | 17.96 | 18.14 | 18.98 |

¹Yields based on early sucrose analysis, assuming that early cane yields are equal to actual yields at harvest.

²LSD for locations means=1.774 t/ha at 10% probability level.

³CV=coefficient of variation.

Table 15.--Indicated yields of 96° sugar, in kilograms per metric ton of cane, from second-ratoon cane on
Lauderhill, Terra Ceia, and Torry muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | Average yield, all farms |
|--------------------------|--|------------------|----------------------|---------------------|--|----------------------|-----------------------|-----------------------------------|
| | Lauderhill muck | | | Terra Ceia muck | | Torry muck | Pompano fine sand | |
| | New Farm, Inc. ¹ 10/30/83 | Duda 11/14/83 | Wedgworth 12/7/83 | Saunders 1/18/84 | South Florida Industries 1/20/84 | Beardsley 2/12/84 | Lykes Bros. 1/2/84 | |
| CP 77-1776..... | 104.2 | 122.5 | 134.6 | 110.0 | 129.6 | 113.0 | 147.4 | 123.1 |
| CP 77-1049..... | 104.4 | 121.8 | 128.5 | 128.2 | 129.8 | 94.4 | 147.1 | 122.0 |
| CP 77-1720..... | 98.5 | 120.0 | 124.7 | 117.6 | 125.3 | 125.2 | 138.5 | 121.3 |
| CP 63-588..... | 94.6 | 111.7 | 116.9 | 115.8 | 121.2 | 107.9 | 141.1 | 115.6 |
| CP 77-1055..... | 104.2 | 116.0 | 119.4 | 108.0 | 120.8 | 103.7 | 136.8 | 115.6 |
| CP 77-1400..... | 100.8 | 112.2 | 118.3 | 109.9 | 115.5 | 101.2 | 138.7 | 113.8 |
| CP 77-1148..... | 100.0 | 105.1 | 111.0 | 109.6 | 124.0 | 106.5 | 136.0 | 113.2 |
| CP 77-1125..... | 94.0 | 103.5 | 113.1 | 122.3 | 120.6 | 102.3 | 130.2 | 112.3 |
| CP 77-1404..... | 94.5 | 107.4 | 112.5 | 111.7 | 113.7 | 112.6 | 132.5 | 112.1 |
| CP 77-1446..... | 109.7 | 109.8 | 121.2 | 104.6 | 104.4 | 91.6 | 126.8 | 109.7 |
| CP 77-1414..... | 98.0 | 108.0 | 109.5 | 104.6 | 114.3 | 103.5 | 122.8 | 108.7 |
| CP 77-1008..... | 86.6 | 98.5 | 116.6 | 99.3 | 107.8 | 96.4 | 123.8 | 104.2 |
| Mean ² | 99.1 | 111.4 | 118.9 | 111.8 | 118.9 | 104.8 | 135.1 | 114.3 |
| LSD: | | | | | | | | |
| 5% level..... | 12.4 | 9.6 | 11.5 | 7.0 | 10.9 | 11.4 | 6.3 | 6.1 |
| 10% level..... | 10.3 | 8.0 | 9.6 | 5.8 | 9.1 | 9.5 | 5.3 | 5.2 |
| CV ³ (%)..... | 8.69 | 5.99 | 6.16 | 4.34 | 6.36 | 7.57 | 3.24 | 6.05 |

¹Polado applied before harvest.

²LSD for location means=2.5 kg/t of cane at 10% probability level.

³CV=coefficient of variation.

Table 16.--Indicated yields of 96° sugar, in metric tons per hectare, from second-ratoon cane on Lauderdale, Terra Ceia, and Torrey muck and on Pompano fine sand

| Clone | Average yield by soil series, farm, and harvest date | | | | | | | Average yield, all farms |
|--------------------------|--|------------------|----------------------|---------------------|-------------------------------------|----------------------|-----------------------|--------------------------|
| | Lauderdale muck | | | Terra Ceia muck | | Torrey muck | Pompano fine sand | |
| | New Farm, Inc. 10/30/83 | Duda 11/14/83 | Wedgworth 12/7/83 | Saunders 1/18/84 | South Florida Industries 1/20/84 | Beardsley 2/12/84 | Lykes Bros. 1/2/84 | |
| CP 77-1414..... | 8.684 | 13.254 | 13.535 | 9.165 | 15.341 | 13.579 | 12.883 | 12.349 |
| CP 77-1055..... | 6.863 | 11.538 | 12.505 | 6.597 | 13.819 | 15.531 | 13.340 | 11.456 |
| CP 77-1404..... | 5.954 | 9.376 | 8.285 | 6.425 | 10.592 | 17.688 | 9.745 | 9.724 |
| CP 77-1125..... | 6.410 | 9.888 | 9.714 | 9.381 | 13.942 | 13.767 | 4.753 | 9.694 |
| CP 77-1008..... | 5.166 | 8.268 | 10.888 | 8.120 | 10.936 | 12.140 | 11.028 | 9.507 |
| CP 63-588..... | 3.691 | 7.739 | 7.406 | 6.843 | 11.614 | 17.105 | 12.111 | 9.501 |
| CP 77-1148..... | 5.235 | 10.201 | 11.119 | 6.714 | 13.772 | 9.618 | 9.732 | 9.484 |
| CP 77-1049..... | 6.571 | 7.886 | 10.327 | 9.825 | 9.512 | 11.459 | 9.562 | 9.306 |
| CP 77-1720..... | 4.774 | 6.850 | 7.798 | 7.381 | 12.935 | 9.864 | 13.793 | 9.056 |
| CP 77-1776..... | 5.230 | 8.728 | 10.043 | 5.442 | 19.422 | 10.652 | 10.479 | 8.570 |
| CP 77-1446..... | 4.978 | 9.565 | 11.214 | 6.210 | 8.873 | 4.783 | 10.933 | 8.080 |
| CP 77-1400..... | 3.481 | 7.594 | 9.515 | 5.306 | 9.252 | 9.903 | 8.819 | 7.696 |
| Mean ¹ | 5.586 | 9.241 | 10.196 | 7.284 | 11.668 | 12.174 | 10.603 | 9.536 |
| LSD: | | | | | | | | |
| 5% level..... | 3.081 | 1.754 | 2.199 | 2.200 | 3.067 | 3.211 | 2.614 | 2.104 |
| 10% level..... | 2.563 | 1.459 | 1.830 | 1.830 | 2.551 | 2.671 | 2.175 | 1.785 |
| CV ² (%)..... | 31.11 | 13.16 | 14.65 | 20.94 | 18.22 | 14.52 | 15.11 | 17.41 |

¹LSD for location means=0.867 t/ha at 10% probability level.
²CV=coefficient of variation.

United States Department of Agriculture
Agricultural Research Service
Beltsville Agricultural Research Center-West
Beltsville, Maryland 20705

OFFICIAL BUSINESS

Penalty for Private Use, \$300



Postage and Fees Paid
U.S. Department of Agriculture
AGR-101